

**DATASHEET**

**Flag-tag Mouse Monoclonal Antibody(C783)**

CAT. NO. AMA00395

**KEY FEATURES**

Target	Flag-tag	Source / Host	Mouse
Reactivity		Clonality	Monoclonal
Applications	WB, IF/ICC, IP	Conjugation	Unconjugated
Form / Buffer	Liquid in PBS, pH 7.4, containing 50% glycerol, 0.05% BSA and 0.01% sodium azide.	Storage	at-20°C

**BACKGROUND**

The FLAG-tag is a short, hydrophilic 8-amino-acid peptide (DYKDDDDK) developed by Sigma-Aldrich for affinity purification and detection of recombinant fusion proteins. FLAG is one of the most popular tags due to its small size, high specificity of available antibodies, and gentle elution conditions for purified proteins. Anti-FLAG antibodies are widely used in WB, IP, IF, IHC, and FACS.

**APPLICATION**

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:2000 - 1:5000
IF/ICC	1:200 - 1:500
IP	1:100 - 1:200

\*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

**PRODUCT OVERVIEW**

Description	Recombinant mouse monoclonal antibody to FLAG-tag
Specificity	Recognizes C-terminal, internal, and N-terminal FLAG-tag fusion proteins
Antibody Type	Primary antibody, Recombinant, Tag
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence of FLAG-tag. The exact sequence is proprietary.
Purification	This antibody is purified by protein A affinity chromatography.
Form/Buffer	Liquid in PBS, pH 7.4, containing 50% glycerol, 0.05% BSA and 0.01% sodium azide.

\*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact [info@arexbio.com](mailto:info@arexbio.com) or your local distributor.


\*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

**DATASHEET**


**Flag-tag Mouse Monoclonal Antibody(C783)**

CAT. NO. AMA00395

**DATA**



Western blot analysis of Flag-tag in 293F transfected with a vector overexpressing Flag tag (A), 293F transfected with an empty vector (B).



Immunofluorescent analysis of FLAG-tag staining in 293T cells transfected with a Flag-tag protein. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AREX® Fluor 488 - conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AREX® Fluor 594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

**STORAGE**

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

**NOTE**

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.