

DATASHEET

VSV-G-tag Mouse Monoclonal Antibody(C2068)

CAT. NO. AMA01680

KEY FEATURES

Target	VSV-G-tag	Source / Host	Mouse
Reactivity		Clonality	Monoclonal
Applications	WB, IF/ICC, IP	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.	Storage	at-20°C

BACKGROUND

The VSV-G-tag is an 11-amino-acid epitope (YTDIEMNRLGK) from the C-terminus of the vesicular stomatitis virus glycoprotein G. It is recognized by the P5D4 monoclonal antibody and used as a fusion tag for the detection, immunoprecipitation, and purification of recombinant proteins.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:2000 - 1:5000
IF/ICC	1:200 - 1:500
IP	1:100 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Mouse monoclonal antibody to VSV-G-tag
Specificity	Recognizes C-terminal, internal, and N-terminal VSV-G-tag fusion proteins.
Antibody Type	Primary antibody, Tag
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence of VSV-G-tag. The exact sequence is proprietary.
Purification	Affinity chromatography
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


VSV-G-tag Mouse Monoclonal Antibody(C2068)

CAT. NO. AMA01680


DATA



Western blot analysis of over-expressed VSV-G-tagged protein in 293T cell lysate.



Immunofluorescent analysis of VSV-G-tag staining in 293T cells transfected with a VSV-G-tag protein. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



Immunoprecipitation of VSV-G-tagged protein from HEK293T cells transfected with vector overexpressing VSV-G tag, using Anti-VSV-G-tag Antibody.

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.