

DATASHEET

ACLY Mouse Monoclonal Antibody(C2487)

CAT. NO. AMA02099

KEY FEATURES

Target	ACLY	Source / Host	Mouse
Reactivity	Human, Mouse, Rat, Monkey	Clonality	Monoclonal
Applications	WB, IHC, IF/ICC, FC	Conjugation	Unconjugated
Form / Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.	Storage	at-20°C

BACKGROUND

Catalyzes the cleavage of citrate into oxaloacetate and acetyl-CoA, the latter serving as common substrate in multiple biochemical reactions in protein, carbohydrate and lipid metabolism.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:100 - 1:500
IF/ICC	1:50 - 1:100
FC	1:100 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Mouse monoclonal to ACLY
Specificity	Recognizes endogenous levels of ACLY protein
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human ACLY expressed in E. Coli
Purification	This antibody is purified through a protein G column.
Molecular Weight	Predicted: 125 kD; Observed: 125 kD kD
Form/Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	ATP-citrate synthase; ATP-citrate (pro-S-)-lyase; ACL; Citrate cleavage enzyme
Gene Symbol	ACLY
Entrez Gene	47(Human); 104112(Mouse); 24159(Rat)
SwissProt	P53396(Human); Q91V92(Mouse); P16638(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


ACLY Mouse Monoclonal Antibody(C2487)

CAT. NO. AMA02099


DATA



Western blot analysis of ACLY expression in HeLa (A), NIH3T3 (B), C6 (C), COS7 (D), Raji (E) whole cell lysates. (Predicted band size: 125 kD; Observed band size: 125 kD)




Immunohistochemical analysis of ACLY staining in human esophageal cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of ACLY staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AREX® Fluor 488 -conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AREX® Fluor 594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

DATASHEET**ACLY Mouse Monoclonal Antibody(C2487)**

CAT. NO. AMA02099

DATA (CONTINUED)

Flow cytometric analysis of HeLa cells using Anti-ACLY Antibody (green) and negative control (red).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.