

DATASHEET

CD99 Mouse Monoclonal Antibody(C2615)

CAT. NO. AMA02227

KEY FEATURES

Target	CD99	Source / Host	Mouse
Reactivity	Human, Mouse	Clonality	Monoclonal
Applications	WB, IHC, FC	Conjugation	Unconjugated
Form / Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.	Storage	at-20°C

BACKGROUND

Involved in T-cell adhesion processes and in spontaneous rosette formation with erythrocytes. Plays a role in a late step of leukocyte extravasation helping leukocytes to overcome the endothelial basement membrane. Acts at the same site as, but independently of, PECAM1. Involved in T-cell adhesion processes .

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:100 - 1:500
FC	1:100 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Mouse monoclonal to CD99
Specificity	Recognizes endogenous levels of CD99 protein
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human CD99 expressed in E. Coli
Purification	This antibody is purified through a protein G column.
Molecular Weight	Predicted: 19 kD; Observed: 32 kD
Form/Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	MIC,2; MIC,2X; MIC,2Y; CD99 antigen; 12E7; E2 antigen; Protein MIC,2; T-cell surface glycoprotein E2; CD99
Gene Symbol	CD99
Entrez Gene	4267(Human)
SwissProt	P14209(Human); Q8VCN6(Mouse)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


CD99 Mouse Monoclonal Antibody(C2615)

CAT. NO. AMA02227


DATA



Western blot analysis of CD99 expression in THP1 (A), U937 (B), MOLT4 (C), L1210 (D) whole cell lysates. (Predicted band size: 19 kD; Observed band size: 32 kD)



Immunohistochemical analysis of CD99 staining in human bladder cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Flow cytometric analysis of Jurkat cells using Anti-CD99 Antibody (green) and negative control (red).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.