

DATASHEET

Beta-catenin Mouse Monoclonal Antibody(C2652)

CAT. NO. AMA02264

KEY FEATURES

Target	Beta-catenin	Source / Host	Mouse
Reactivity	Human, Mouse, Rat, Monkey	Clonality	Monoclonal
Applications	WB, IHC, FC	Conjugation	Unconjugated
Form / Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.	Storage	at-20°C

BACKGROUND

Key downstream component of the canonical Wnt signaling pathway . In the absence of Wnt, forms a complex with AXIN1, AXIN2, APC, CSNK1A1 and GSK3B that promotes phosphorylation on N-terminal Ser and Thr residues and ubiquitination of CTNNB1 via BTRC and its subsequent degradation by the proteasome . In the presence of Wnt ligand, CTNNB1 is not ubiquitinated and accumulates in the nucleus, where it acts as a coactivator for transcription factors of the TCF/LEF family, leading to activate Wnt responsive genes . Also acts as a coactivator for other transcription factors, such as NR5A2 . Promotes epithelial to mesenchymal transition/mesenchymal to epithelial transition (EMT/MET) via driving transcription of CTNNB1/TCF-target genes .

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:100 - 1:500
FC	1:100 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Mouse monoclonal to Beta-catenin
Specificity	Recognizes endogenous levels of Beta-catenin protein
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human Beta-catenin expressed in E. Coli
Purification	This antibody is purified through a protein G column.
Molecular Weight	Predicted: 86 kD; Observed: 86 kD
Form/Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	CTNNB; Catenin beta-1; Beta-catenin
Gene Symbol	CTNNB1
Entrez Gene	1499(Human); 12387(Mouse); 84353(Rat)
SwissProt	P35222(Human); Q02248(Mouse); Q9WU82(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


Beta-catenin Mouse Monoclonal Antibody(C2652)

CAT. NO. AMA02264


DATA



Western blot analysis of Beta-catenin expression in HeLa (A), HepG2 (B), NIH3T3 (C), MCF7 (D), C6 (E), COS7 (F), K562 (G), Jurkat (H), A549 (I), SHSY5Y (J), BEL7402 (K), HEK293 (L), HEK293-6e (M) whole cell lysates. (Predicted band size: 86 kD; Observed band size: 86 kD)



Immunohistochemical analysis of Beta-catenin staining in human cervical cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Flow cytometric analysis of HepG2 cells using Anti-Beta-catenin Antibody (green) and negative control (red).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.