

DATASHEET

DCTN4 Mouse Monoclonal Antibody(C2658)

CAT. NO. AMA02270

KEY FEATURES

Target	DCTN4	Source / Host	Mouse
Reactivity	Human	Clonality	Monoclonal
Applications	WB, IHC, IF/ICC, FC	Conjugation	Unconjugated
Form / Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.	Storage	at-20°C

BACKGROUND

Part of the dynactin complex that activates the molecular motor dynein for ultra-processive transport along microtubules.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:100 - 1:500
IF/ICC	1:100 - 1:500
FC	1:100 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Mouse monoclonal to DCTN4
Specificity	Recognizes endogenous levels of DCTN4 protein
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human DCTN4 expressed in E. Coli
Purification	This antibody is purified through a protein G column.
Molecular Weight	Predicted: 52 kD; Observed: 54 kD
Form/Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	Dynactin subunit 4; Dyn4; Dynactin subunit p62
Gene Symbol	DCTN4
Entrez Gene	51164(Human)
SwissProt	Q9UJW0(Human)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


DCTN4 Mouse Monoclonal Antibody(C2658)

CAT. NO. AMA02270


DATA



Western blot analysis of DCTN4 expression in Raw264.7 (A), NIH3T3 (B) whole cell lysates. (Predicted band size: 52 kD; Observed band size: 54 kD)




Immunohistochemical analysis of DCTN4 staining in human cervical cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of DCTN4 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AREX® Fluor 488 -conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AREX® Fluor 594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

DATASHEET**DCTN4 Mouse Monoclonal Antibody(C2658)**

CAT. NO. AMA02270

DATA (CONTINUED)

Flow cytometric analysis of A549 cells using Anti-DCTN4 Antibody (green) and negative control (red).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.