

DATASHEET

EIF4B Mouse Monoclonal Antibody(C2681)

CAT. NO. AMA02293

KEY FEATURES

Target	EIF4B	Source / Host	Mouse
Reactivity	Human, Mouse	Clonality	Monoclonal
Applications	WB, IHC, IF/ICC, FC	Conjugation	Unconjugated
Form / Buffer	Mouse IgG2b. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.	Storage	at-20°C

BACKGROUND

Required for the binding of mRNA to ribosomes. Functions in close association with EIF4-F and EIF4-A. Binds near the 5'-terminal cap of mRNA in presence of EIF-4F and ATP. Promotes the ATPase activity and the ATP-dependent RNA unwinding activity of both EIF4-A and EIF4-F.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:100 - 1:500
IF/ICC	1:100 - 1:500
FC	1:100 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Mouse monoclonal to EIF4B
Specificity	Recognizes endogenous levels of EIF4B protein
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human EIF4B expressed in E. Coli
Purification	This antibody is purified through a protein G column.
Molecular Weight	Predicted: 80 kD; Observed: 80 kD kD
Form/Buffer	Mouse IgG2b. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	EukaryotIC, translation initiation factor 4B; eIF-4B
Gene Symbol	EIF4B
Entrez Gene	1975(Human)
SwissProt	P23588(Human); Q8BGD9(Mouse)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


EIF4B Mouse Monoclonal Antibody(C2681)

CAT. NO. AMA02293


DATA



Western blot analysis of EIF4B expression in A549 (A), A431 (B), HepG2 (C), HEK293 (D), HeLa (E), Jurkat (F), K562 (G), NIH3T3 (H), MCF7 (I) whole cell lysates. (Predicted band size: 80 kD; Observed band size: 80 kD)




Immunohistochemical analysis of EIF4B staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of EIF4B staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AREX® Fluor 488 -conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AREX® Fluor 594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

DATASHEET**EIF4B Mouse Monoclonal Antibody(C2681)**

CAT. NO. AMA02293

DATA (CONTINUED)

Flow cytometric analysis of HeLa cells using Anti-EIF4B Antibody (green) and negative control (red).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.