

DATASHEET

hnRNP M Mouse Monoclonal Antibody(C2748)

CAT. NO. AMA02360

KEY FEATURES

Target	hnRNP M	Source / Host	Mouse
Reactivity	Human, Mouse, Rat	Clonality	Monoclonal
Applications	WB, IHC, IF/ICC, FC	Conjugation	Unconjugated
Form / Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.	Storage	at-20°C

BACKGROUND

Pre-mRNA binding protein in vivo, binds avidly to poly(G) and poly(U) RNA homopolymers in vitro. Involved in splicing. Acts as a receptor for carcinoembryonic antigen in Kupffer cells, may initiate a series of signaling events leading to tyrosine phosphorylation of proteins and induction of IL-1 alpha, IL-6, IL-10 and tumor necrosis factor alpha cytokines.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:100 - 1:500
IF/ICC	1:100 - 1:500
FC	1:100 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Mouse monoclonal to hnRNP M
Specificity	Recognizes endogenous levels of hnRNP M protein
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human hnRNP M expressed in E. Coli
Purification	This antibody is purified through a protein G column.
Molecular Weight	Predicted: 78 kD; Observed: 80 kD
Form/Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	HNRPM; NAGR1; Heterogeneous nuclear ribonucleoprotein M; hnRNP M
Gene Symbol	HNRNPM
Entrez Gene	4670(Human); 76936(Mouse); 116655(Rat)
SwissProt	P52272(Human); Q9D0E1(Mouse); Q62826(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


hnRNP M Mouse Monoclonal Antibody(C2748)

CAT. NO. AMA02360


DATA



Western blot analysis of hnRNP M expression in Raji (A), HeLa (B), NIH/3T3 (C), A431 (D), A549 (E), HepG2 (F), PC12 (G), U251 (H) whole cell lysates. (Predicted band size: 78 kD; Observed band size: 80 kD)




Immunohistochemical analysis of hnRNP M staining in human esophageal cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of hnRNP M staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AREX® Fluor 488 -conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AREX® Fluor 594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

DATASHEET**hnRNP M Mouse Monoclonal Antibody(C2748)**

CAT. NO. AMA02360

DATA (CONTINUED)

Flow cytometric analysis of HL60 cells using Anti-hnRNP M Antibody (green) and negative control (red).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.