

DATASHEET

CD138 Mouse Monoclonal Antibody(C2951)

CAT. NO. AMA02563

KEY FEATURES

Target	CD138	Source / Host	Mouse
Reactivity	Human, Mouse	Clonality	Monoclonal
Applications	WB, IHC, IF/ICC, FC	Conjugation	Unconjugated
Form / Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.	Storage	at-20°C

BACKGROUND

Cell surface proteoglycan that contains both heparan sulfate and chondroitin sulfate and that links the cytoskeleton to the interstitial matrix . Regulates exosome biogenesis in concert with SDCBP and PDCD6IP . Able to induce its own expression in dental mesenchymal cells and also in the neighboring dental epithelial cells via an MSX1-mediated pathway .

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:100 - 1:500
IF/ICC	1:100 - 1:500
FC	1:100 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Mouse monoclonal to CD138
Specificity	Recognizes endogenous levels of CD138 protein
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human CD138 expressed in E. Coli
Purification	This antibody is purified through a protein G column.
Molecular Weight	Predicted: 33 kD; Observed: 84 kD kD
Form/Buffer	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	SDC; Syndecan-1; SYND1; CD138
Gene Symbol	SDC1
Entrez Gene	6382(Human); 25216(Rat)
SwissProt	P18827(Human); P18828(Mouse)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


CD138 Mouse Monoclonal Antibody(C2951)

CAT. NO. AMA02563


DATA



Western blot analysis of CD138 expression in MCF7 (A), HeLa (B), HepG2 (C), T47D (D), SW620 (E), Jurkat (F), NIH/3T3 (G) whole cell lysates. (Predicted band size: 33 kD; Observed band size: 84 kD)




Immunohistochemical analysis of CD138 staining in human ovarian cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of CD138 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AREX® Fluor 488 -conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AREX® Fluor 594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

DATASHEET**CD138 Mouse Monoclonal Antibody(C2951)**

CAT. NO. AMA02563

DATA (CONTINUED)

Flow cytometric analysis of HepG2 cells using Anti-CD138 Antibody (green) and negative control (red).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.