

DATASHEET

PCNA Mouse Monoclonal Antibody(C3481)

CAT. NO. AMA03093

KEY FEATURES

Target	PCNA	Source / Host	Mouse
Reactivity	Human, Mouse, Rat, Monkey, Hamster	Clonality	Monoclonal
Applications	WB, IF/ICC, IP	Conjugation	Unconjugated
Form / Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.	Storage	at-20°C

BACKGROUND

Confers DNA tethering and processivity to DNA polymerases and other proteins. Auxiliary protein of DNA polymerase delta and epsilon, is involved in the control of DNA replication by increasing the polymerases' processivity during elongation of the leading strand. Induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic-aprimidinic (AP) endonuclease, APEX2 activities. Has to be loaded onto DNA in order to be able to stimulate APEX2. Plays a key role in DNA damage response (DDR) by being conveniently positioned at the replication fork to coordinate DNA replication with DNA repair and DNA damage tolerance pathways.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IF/ICC	1:50 - 1:100
IP	1:10 - 1:50

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Mouse monoclonal antibody to PCNA
Specificity	Recognizes endogenous levels of PCNA protein.
Antibody Type	Primary antibody
Immunogen	Purified recombinant human PCNA protein fragments expressed in E.coli.
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 29 kD; Observed: 36 kD
Form/Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Alternative Names	Proliferating cell nuclear antigen; PCNA; Cyclin
Gene Symbol	PCNA
Entrez Gene	5111(Human); 18538(Mouse); 25737(Rat)
SwissProt	P12004(Human); P17918(Mouse); P04961(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


PCNA Mouse Monoclonal Antibody(C3481)

CAT. NO. AMA03093

DATA



Western blot analysis of PCNA expression in HeLa (A), NIH3T3 (B), COS7 (C), C6 (D), CHOK1 (E) whole cell lysates. (Predicted band size: 29 kD; Observed band size: 36 kD)



Immunofluorescent analysis of PCNA staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AREX® Fluor 488 -conjugated secondary antibody (green) in PBS at room temperature in the dark.

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.