

DATASHEET

Histamine H2 Receptor Rabbit Polyclonal Antibody

CAT. NO. APA07097

KEY FEATURES

Target	Histamine H2 Receptor	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at-20°C

BACKGROUND

G-protein coupled receptor for histamine, primarily mediating gastric acid secretion. Predominantly expressed in the gastric mucosa, couples to G(s) G alpha proteins upon histamine binding, leading to activation of adenylate cyclase and increased intracellular cyclic AMP (cAMP) levels. G alpha proteins upon histamine binding, leading to activation of adenylate cyclase and increased intracellular cyclic AMP (cAMP) levels. This signaling cascade stimulates parietal cells to secrete hydrochloric acid, playing a key role in digestive physiology. Also expressed in other tissues, including the heart and central nervous system, where it may contribute to cardiac stimulation and modulate neurotransmitter release.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IF/ICC	1:100 - 1:500

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to Histamine H2 Receptor
Specificity	Recognizes endogenous levels of Histamine H2 Receptor protein.
Antibody Type	Primary antibody
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Histamine H2 Receptor. The exact sequence is proprietary.
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 40 kD; Observed: 40 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	Histamine H2 receptor; H2R; HH2R; Gastric receptor I
Gene Symbol	HRH2
Entrez Gene	3274(Human); 15466(Mouse); 25461(Rat)
SwissProt	P25021(Human); P97292(Mouse); P25102(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


Histamine H2 Receptor Rabbit Polyclonal Antibody

CAT. NO. APA07097

DATA



Western blot analysis of Histamine H2 Receptor expression in mouse kidney (A), rat liver (B) whole cell lysates. (Predicted band size: 40 kD; Observed band size: 40 kD)



Immunofluorescent analysis of Histamine H2 Receptor staining in COLO205 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.