

DATASHEET

TRIM59 Rabbit Polyclonal Antibody

CAT. NO. APA10647

KEY FEATURES

Target	TRIM59	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine	Clonality	Polyclonal
Applications	WB, IHC, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

E3 ubiquitin ligase involved in different processes such as development and immune response . Serves as a negative regulator for innate immune signaling pathways by suppressing RLR-induced activation of IRF3/7 and NF-kappa-B via interaction with adapter ECSIT . Regulates autophagy through modulating both the transcription and the ubiquitination of BECN1 . On the one hand, regulates the transcription of BECN1 through negatively modulating the NF-kappa-B pathway. On the other hand, regulates TRAF6-mediated 'Lys-63'-linked ubiquitination of BECN1, thus affecting the formation of the BECN1-PIK3C3 complex. In addition, mediates 'Lys-48'-linked ubiquitination of TRAF6 and thereby promotes TRAF6 proteasomal degradation .

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:50 - 1:100
IF/ICC	1:100 - 1:500

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to TRIM59
Specificity	Recognizes endogenous levels of TRIM59 protein.
Antibody Type	Primary antibody
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TRIM59. The exact sequence is proprietary.
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 47 kD; Observed: 50 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	RNF104; TRIM57; TSBF1; Tripartite motif-containing protein 59; RING finger protein 104; Tumor suppressor TSBF-1
Gene Symbol	TRIM59
Entrez Gene	286827(Human); 66949(Mouse)
SwissProt	Q8IWR1(Human); Q922Y2(Mouse)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


TRIM59 Rabbit Polyclonal Antibody

CAT. NO. APA10647


DATA



Western blot analysis of TRIM59 expression in Jurkat (A), HEK293T (B), rat brain (C), mouse brain (D) whole cell lysates. (Predicted band size: 47 kD; Observed band size: 50 kD)



Immunohistochemical analysis of TRIM59 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of TRIM59 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with Alexa Fluor 647-conjugated secondary antibody (red) in PBS at room temperature in the dark.

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.