

DATASHEET

Synaptotagmin (Phospho-T202/199) Rabbit Polyclonal Antibody

CAT. NO. APA10906

KEY FEATURES

Target	Synaptotagmin (Phospho-T202/199)	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB, IHC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

Calcium sensor that participates in triggering neurotransmitter release at the synapse . May have a regulatory role in the membrane interactions during trafficking of synaptic vesicles at the active zone of the synapse . It binds acidic phospholipids with a specificity that requires the presence of both an acidic head group and a diacyl backbone. A Ca(2+)-dependent interaction between synaptotagmin and putative receptors for activated protein kinase C has also been reported. It can bind to at least three additional proteins in a Ca(2+)-independent manner; these are neuroligins, syntaxin and AP2. Plays a role in dendrite formation by melanocytes -dependent interaction between synaptotagmin and putative receptors for activated protein kinase C has also been reported.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:50 - 1:100

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to Synaptotagmin (Phospho-T202/199)
Specificity	Recognizes endogenous levels of Synaptotagmin protein only when phosphorylated at T202/199.
Antibody Type	Primary antibody, Tag
Immunogen	KLH-conjugated synthetic phosphopeptide corresponding to residues surrounding T202/199 of human Synaptotagmin protein. The exact sequence is proprietary.
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 47; Observed: 47 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	SYT1; SVP65; SYT; Synaptotagmin-1; Synaptotagmin I; SytI; p65; SYT2; Synaptotagmin-2; Synaptotagmin II; SytII
Gene Symbol	SYT1; SYT2
Entrez Gene	6857; 127833(Human); 20979; 20980(Mouse); 25716; 24805(Rat)
SwissProt	P21579; Q8N9I0(Human); P46096; P46097(Mouse); P21707; P29101(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


Synaptotagmin (Phospho-T202/199) Rabbit Polyclonal Antibody

CAT. NO. APA10906

DATA



Western blot analysis of Synaptotagmin (Phospho-T202/199) expression in K562 (A), HEK293T (B), U87MG (C), A549 (D) whole cell lysates. (Predicted band size: 47; 46 kD; Observed band size: 47 kD)



Immunohistochemical analysis of Synaptotagmin (Phospho-T202/199) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.