

DATASHEET

YEATS2 Rabbit Polyclonal Antibody

CAT. NO. APA11019

KEY FEATURES

Target	YEATS2	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

Chromatin reader component of the ATAC complex, a complex with histone acetyltransferase activity on histones H3 and H4 . YEATS2 specifically recognizes and binds histone H3 crotonylated at 'Lys-27' (H3K27cr) . Crotonylation marks active promoters and enhancers and confers resistance to transcriptional repressors .

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IF/ICC	1:50 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to YEATS2
Specificity	Recognizes endogenous levels of YEATS2 protein.
Antibody Type	Primary antibody
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human YEATS2. The exact sequence is proprietary.
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 150 kD; Observed: 150 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	KIAA1197; YEATS domain-containing protein 2
Gene Symbol	YEATS2
Entrez Gene	55689(Human); 208146(Mouse)
SwissProt	Q9ULM3(Human); Q3TUF7(Mouse)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


YEATS2 Rabbit Polyclonal Antibody

CAT. NO. APA11019

DATA



Western blot analysis of YEATS2 expression in C6 (A), SP20 (B), A375 (C), HeLa (D) whole cell lysates. (Predicted band size: 150 kD; Observed band size: 150 kD)



Immunofluorescent analysis of YEATS2 staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AREX® Fluor 488 -conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AREX® Fluor 594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.