

DATASHEET

Claudin 7 (Phospho-Y210) Rabbit Polyclonal Antibody

CAT. NO. APA11674

KEY FEATURES

Target	Claudin 7 (Phospho-Y210)	Source / Host	Rabbit
Reactivity	Human, Rat	Clonality	Polyclonal
Applications	WB, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

Plays a major role in tight junction-specific obliteration of the intercellular space.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IF/ICC	1:50 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to Claudin 7 (Phospho-Y210)
Specificity	Recognizes endogenous levels of Claudin 7 protein only when phosphorylated at Y210.
Antibody Type	Primary antibody
Immunogen	KLH-conjugated synthetic phosphopeptide corresponding to residues surrounding Y210 of human Claudin 7 protein. The exact sequence is proprietary.
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 22 kD; Observed: 27 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	CEPTL2; CPETRL2; Claudin-7; CLDN-7
Gene Symbol	CLDN7
Entrez Gene	1366(Human); 65132(Rat)
SwissProt	O95471(Human); Q9Z1L1(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


Claudin 7 (Phospho-Y210) Rabbit Polyclonal Antibody

CAT. NO. APA11674

DATA



Western blot analysis of Claudin 7 (Phospho-Y210) expression in rat liver (A), rat kidney (B) whole cell lysates. (Predicted band size: 22 kD; Observed band size: 27 kD)



Immunofluorescent analysis of Claudin 7 (Phospho-Y210) staining in MG63 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AREX® Fluor 488 - conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AREX® Fluor 594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.