

DATASHEET

NBC1 Rabbit Polyclonal Antibody

CAT. NO. APA12720

KEY FEATURES

Target	NBC1	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

Electrogenic sodium/bicarbonate cotransporter with a Na(+):HCO3(-) stoichiometry varying from 1:2 to 1:3. May regulate bicarbonate influx/efflux at the basolateral membrane of cells and regulate intracellular pH.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IF/ICC	1:50 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to NBC1
Specificity	Recognizes endogenous levels of NBC1 protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human NBC1
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 72; Observed: 135 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	NBC; NBC1; NBCE1; Electrogenic sodium bicarbonate cotransporter 1; Sodium bicarbonate cotransporter; Na(+)/HCO3(-) cotransporter; Solute carrier family 4 member 4; kNBC1
Gene Symbol	SLC4A4
Entrez Gene	8671(Human); 54403(Mouse); 84484(Rat)
SwissProt	Q9Y6R1(Human); O88343(Mouse); Q9JI66(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


NBC1 Rabbit Polyclonal Antibody

CAT. NO. APA12720

DATA



Western blot analysis of NBC1 expression in PC3 (A), A549 (B) whole cell lysates. (Predicted band size: 72; 112; 116; 121; 123 kD; Observed band size: 135 kD)



Immunofluorescent analysis of NBC1 staining in C6 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AREX® Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.