

DATASHEET

Neurokinin B Rabbit Polyclonal Antibody

CAT. NO. APA12857

KEY FEATURES

Target	Neurokinin B	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at-20°C

BACKGROUND

Is a ligand for TACR3, and triggers G protein-coupled receptor signaling via activation of G(q) and phosphatidylinositol hydrolysis by phospholipase C and phosphatidylinositol hydrolysis by phospholipase C . Binding to TACR3 also triggers signaling via activation of adenylate cyclase activity which results in increased intracellular levels of cyclic AMP (cAMP) . Is a critical central regulator of gonadal function .

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:2000
IF/ICC	1:10 - 1:100

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to Neurokinin B
Specificity	Recognizes endogenous levels of Neurokinin B protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human Neurokinin B
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 11; Observed: 16 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	NKNB; Tachykinin-3; ZNEUROK1
Gene Symbol	TAC3
Entrez Gene	6866(Human); 21334(Mouse); 29191(Rat)
SwissProt	Q9UHF0(Human); P55099(Mouse); P08435(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


Neurokinin B Rabbit Polyclonal Antibody

CAT. NO. APA12857

DATA



Western blot analysis of Neurokinin B expression in Jurkat (A), mouse spleen (B), rat kidney (C), rat brain (D) whole cell lysates. (Predicted band size: 11; 13; 14 kD; Observed band size: 16 kD)



Immunofluorescent analysis of Neurokinin B staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.