

DATASHEET

GDA Rabbit Polyclonal Antibody

CAT. NO. APA12901

KEY FEATURES

Target	GDA	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

Catalyzes the hydrolytic deamination of guanine, producing xanthine and ammonia.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
----	----------------

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW


Description	Rabbit polyclonal antibody to GDA
Specificity	Recognizes endogenous levels of GDA protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human GDA
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 42; Observed: 51 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	KIAA1258; Guanine deaminase; Guanase; Guanine aminase; Guanine aminohydrolase; GAH; p51-nedasin
Gene Symbol	GDA
Entrez Gene	9615(Human)
SwissProt	Q9Y2T3(Human)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.

*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET**GDA Rabbit Polyclonal Antibody**

CAT. NO. APA12901

DATA

Western blot analysis of GDA expression in HeLa (A), SKOV3 (B), mouse lung (C), rat brain (D) whole cell lysates. (Predicted band size: 42; 51; 52 kD; Observed band size: 51 kD)

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.