

**DATASHEET**

**MSRB3 Rabbit Polyclonal Antibody**

CAT. NO. APA13210

**KEY FEATURES**

Target	MSRB3	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

**BACKGROUND**

Catalyzes the reduction of free and protein-bound methionine sulfoxide to methionine. Isoform 2 is essential for hearing.

**APPLICATION**

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:2000
IF/ICC	1:50 - 1:200

\*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

**PRODUCT OVERVIEW**

Description	Rabbit polyclonal antibody to MSRB3
Specificity	Recognizes endogenous levels of MSRB3 protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human MSRB3
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 20 kD; Observed: 25 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	Methionine-R-sulfoxide reductase B3; MsrB3
Gene Symbol	MSRB3
Entrez Gene	253827(Human); 320183(Mouse)
SwissProt	Q8IXL7(Human); Q8BU85(Mouse)

\*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact [info@arexbio.com](mailto:info@arexbio.com) or your local distributor.


\*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

**DATASHEET**


**MSRB3 Rabbit Polyclonal Antibody**

CAT. NO. APA13210

**DATA**



Western blot analysis of MSRB3 expression in mouse testis (A), mouse brain (B), rat lung (C) whole cell lysates. (Predicted band size: 20 kD; Observed band size: 25 kD)



Immunofluorescent analysis of MSRB3 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AREX® Fluor 488 -conjugated secondary antibody (green) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

**STORAGE**

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

**NOTE**

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.