

DATASHEET

Unc18-3 Rabbit Polyclonal Antibody

CAT. NO. APA13232

KEY FEATURES

Target	Unc18-3	Source / Host	Rabbit
Reactivity	Human, Mouse	Clonality	Polyclonal
Applications	WB, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

Together with STX4 and VAMP2, may play a role in insulin-dependent movement of GLUT4 and in docking/fusion of intracellular GLUT4-containing vesicles with the cell surface in adipocytes.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:2000
IF/ICC	1:50 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to Unc18-3
Specificity	Recognizes endogenous levels of Unc18-3 protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human Unc18-3
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 67 kD; Observed: 68 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	Syntaxin-binding protein 3; Platelet Sec1 protein; PSP; Protein unc-18 homolog 3; Unc18-3; Protein unc-18 homolog C; Unc-18C
Gene Symbol	STXBP3
Entrez Gene	6814(Human); 20912(Mouse)
SwissProt	O00186(Human); Q60770(Mouse)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


Unc18-3 Rabbit Polyclonal Antibody

CAT. NO. APA13232

DATA



Western blot analysis of Unc18-3 expression in Jurkat (A), mouse kidney (B), mouse brain (C), rat brain (D) whole cell lysates. (Predicted band size: 67 kD; Observed band size: 68 kD)



Immunofluorescent analysis of Unc18-3 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AREX® Fluor 488 -conjugated secondary antibody (green) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.