

DATASHEET

Carbonic Anhydrase 3 Rabbit Polyclonal Antibody

CAT. NO. APA13344

KEY FEATURES

Target	Carbonic Anhydrase 3	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB, IHC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

Reversible hydration of carbon dioxide.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:50 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to Carbonic Anhydrase 3
Specificity	Recognizes endogenous levels of Carbonic Anhydrase 3 protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human Carbonic Anhydrase 3
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 29 kD; Observed: 30 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	Carbonic anhydrase 3; Carbonate dehydratase III; Carbonic anhydrase III; CA-III
Gene Symbol	CA3
Entrez Gene	761(Human); 12350(Mouse); 54232(Rat)
SwissProt	P07451(Human); P16015(Mouse); P14141(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


Carbonic Anhydrase 3 Rabbit Polyclonal Antibody

CAT. NO. APA13344

DATA



Western blot analysis of Carbonic Anhydrase 3 expression in mouse kidney (A), mouse lung (B) whole cell lysates. (Predicted band size: 29 kD; Observed band size: 30 kD)



Immunohistochemical analysis of Carbonic Anhydrase 3 staining in human testis formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.