

DATASHEET

H-FABP Rabbit Polyclonal Antibody

CAT. NO. APA13434

KEY FEATURES

Target	H-FABP	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB, IHC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at-20°C

BACKGROUND

FABPs are thought to play a role in the intracellular transport of long-chain fatty acids and their acyl-CoA esters.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:50 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW


Description	Rabbit polyclonal antibody to H-FABP
Specificity	Recognizes endogenous levels of H-FABP protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human H-FABP
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 14 kD; Observed: 14 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	FABP11; MDGI; Fatty acid-binding protein, heart; Fatty acid-binding protein 3; Heart-type fatty acid-binding protein; H-FABP; Mammary-derived growth inhibitor; MDGI; Muscle fatty acid-binding protein; M-FABP
Gene Symbol	FABP3
Entrez Gene	2170(Human); 14077(Mouse); 79131(Rat)
SwissProt	P05413(Human); P11404(Mouse); P07483(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET**H-FABP Rabbit Polyclonal Antibody**

CAT. NO. APA13434

DATA

Western blot analysis of H-FABP expression in mouse heart (A) whole cell lysates.
(Predicted band size: 14 kD; Observed band size: 14 kD)



Immunohistochemical analysis of H-FABP staining in human testis formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.