

DATASHEET

Nucleostemin Rabbit Polyclonal Antibody

CAT. NO. APA13463

KEY FEATURES

Target	Nucleostemin	Source / Host	Rabbit
Reactivity	Human	Clonality	Polyclonal
Applications	WB, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

May be required to maintain the proliferative capacity of stem cells. Stabilizes MDM2 by preventing its ubiquitination, and hence proteasomal degradation .

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:2000
IF/ICC	1:10 - 1:100

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW


Description	Rabbit polyclonal antibody to Nucleostemin
Specificity	Recognizes endogenous levels of Nucleostemin protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human Nucleostemin
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 60; Observed: 62 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	E2IG3; NS; Guanine nucleotide-binding protein-like 3; E2-induced gene 3 protein; Novel nucleolar protein 47; NNP47; Nucleolar GTP-binding protein 3; Nucleostemin
Gene Symbol	GNL3
Entrez Gene	26354(Human)
SwissProt	Q9BVP2(Human)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET**Nucleostemin Rabbit Polyclonal Antibody**

CAT. NO. APA13463

DATA

Western blot analysis of Nucleostemin expression in HEK293T (A) whole cell lysates. (Predicted band size: 60; 61 kD; Observed band size: 62 kD)



Immunofluorescent analysis of Nucleostemin staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.