

**DATASHEET**

**hnRNP E2 Rabbit Polyclonal Antibody**

CAT. NO. APA13603

**KEY FEATURES**

Target	hnRNP E2	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB, IHC, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

**BACKGROUND**

Single-stranded nucleic acid binding protein that binds preferentially to oligo dC . Major cellular poly(rC)-binding protein . Also binds poly(rU) . Acts as a negative regulator of antiviral signaling . Negatively regulates cellular antiviral responses mediated by MAVS signaling . It acts as an adapter between MAVS and the E3 ubiquitin ligase ITCH, therefore triggering MAVS ubiquitination and degradation . Negatively regulates the cGAS-STING pathway via interaction with CGAS, preventing the formation of liquid-like droplets in which CGAS is activated . Together with PCBP1, required for erythropoiesis, possibly by regulating mRNA splicing .

**APPLICATION**

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:2000
IHC	1:50 - 1:200
IF/ICC	1:50 - 1:200

\*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

**PRODUCT OVERVIEW**

Description	Rabbit polyclonal antibody to hnRNP E2
Specificity	Recognizes endogenous levels of hnRNP E2 protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human hnRNP E2
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 33; Observed: 39 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	Poly(rC)-binding protein 2; Alpha-CP2; Heterogeneous nuclear ribonucleoprotein E2; hnRNP E2
Gene Symbol	PCBP2
Entrez Gene	5094(Human); 18521(Mouse)
SwissProt	Q15366(Human); Q61990(Mouse)

\*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact [info@arexbio.com](mailto:info@arexbio.com) or your local distributor.


\*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

**DATASHEET**


**hnRNP E2 Rabbit Polyclonal Antibody**

CAT. NO. APA13603


**DATA**



Western blot analysis of hnRNP E2 expression in SW620 (A), MCF7 (B), mouse spleen (C) whole cell lysates. (Predicted band size: 33; 34; 35; 38 kD; Observed band size: 39 kD)



Immunohistochemical analysis of hnRNP E2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of hnRNP E2 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

**STORAGE**

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

**NOTE**

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.