DATASHEET

ESAM Rabbit Polyclonal Antibody

CAT. NO. APA14093

KEY FEATURES

Target	ESAM	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.	Storage	at-20°C

BACKGROUND

Can mediate aggregation most likely through a homophilic molecular interaction.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:2000

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to ESAM
Specificity	Recognizes endogenous levels of ESAM protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human ESAM
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 15; Observed: 60 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	Endothelial cell-selective adhesion molecule
Gene Symbol	ESAM
Entrez Gene	90952(Human); 69524(Mouse); 300519(Rat)
SwissProt	Q96AP7(Human); Q925F2(Mouse); Q6AYD4(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET

ESAM Rabbit Polyclonal Antibody

CAT. NO. APA14093

DATA



Western blot analysis of ESAM expression in HT29 (A), HepG2 (B), A549 (C) whole cell lysates. (Predicted band size: 15; 41 kD; Observed band size: 60 kD)

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.