

DATASHEET

PGAM1 Rabbit Polyclonal Antibody

CAT. NO. APA14337

KEY FEATURES

Target	PGAM1	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

Catalyzes the interconversion of 2-phosphoglycerate and 3-phosphoglycerate, a crucial step in glycolysis, by using 2,3-bisphosphoglycerate. Also catalyzes the interconversion of (2R)-2,3-bisphosphoglycerate and (2R)-3-phospho-glyceroyl phosphate.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:2000
IF/ICC	1:50 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to PGAM1
Specificity	Recognizes endogenous levels of PGAM1 protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human PGAM1
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 28 kD; Observed: 29 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	PGAMA; Phosphoglycerate mutase 1; BPG-dependent PGAM 1; Phosphoglycerate mutase isozyme B; PGAM-B
Gene Symbol	PGAM1
Entrez Gene	5223(Human); 18648(Mouse); 24642(Rat)
SwissProt	P18669(Human); Q9DBJ1(Mouse); P25113(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


PGAM1 Rabbit Polyclonal Antibody

CAT. NO. APA14337

DATA



Western blot analysis of PGAM1 expression in MCF7 (A), NIH3T3 (B), mouse brain (C), rat spinal cord (D) whole cell lysates. (Predicted band size: 28 kD; Observed band size: 29 kD)



Immunofluorescent analysis of PGAM1 staining in L929 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AREX® Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.