

DATASHEET

BMF Rabbit Polyclonal Antibody

CAT. NO. APA14475

KEY FEATURES

Target	BMF	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

May play a role in apoptosis. Isoform 1 seems to be the main initiator.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
----	----------------

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW


Description	Rabbit polyclonal antibody to BMF
Specificity	Recognizes endogenous levels of BMF protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human BMF
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 14; Observed: 21 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	Bcl-2-modifying factor
Gene Symbol	BMF
Entrez Gene	90427(Human); 171543(Mouse); 246142(Rat)
SwissProt	Q96LC9(Human); Q91ZE9(Mouse); Q8K589(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.

*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET**BMF Rabbit Polyclonal Antibody**

CAT. NO. APA14475

DATA

Western blot analysis of BMF expression in SW480 (A), mouse lung (B), rat liver (C) whole cell lysates. (Predicted band size: 14; 17; 20 kD; Observed band size: 21 kD)

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.