

DATASHEET

KIFC3 Rabbit Polyclonal Antibody

CAT. NO. APA14829

KEY FEATURES

Target	KIFC3	Source / Host	Rabbit
Reactivity	Human, Rat	Clonality	Polyclonal
Applications	WB, IP	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at-20°C

BACKGROUND

Minus-end microtubule-dependent motor protein. Involved in apically targeted transport . Required for zonula adherens maintenance.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:2000
IP	1:50 - 1:100

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW


Description	Rabbit polyclonal antibody to KIFC3
Specificity	Recognizes endogenous levels of KIFC3 protein.
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human KIFC3
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 92 kD; Observed: 120 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	Kinesin-like protein KIFC3
Gene Symbol	KIFC3
Entrez Gene	3801(Human)
SwissProt	Q9BVG8(Human)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.

*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET**KIFC3 Rabbit Polyclonal Antibody**

CAT. NO. APA14829

DATA

Western blot analysis of KIFC3 expression in U87 (A), rat kidney (B) whole cell lysates. (Predicted band size: 92 kD; Observed band size: 120 kD)

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.