

DATASHEET

Septin 5 Rabbit Polyclonal Antibody

CAT. NO. APA14985

KEY FEATURES

Target	Septin 5	Source / Host	Rabbit
Reactivity	Human, Mouse, Rat	Clonality	Polyclonal
Applications	WB, IHC, IF/ICC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

Filament-forming cytoskeletal GTPase . May play a role in cytokinesis (Potential). May play a role in platelet secretion .

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:2000
IHC	1:50 - 1:200
IF/ICC	1:50 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW

Description	Rabbit polyclonal antibody to Septin 5
Specificity	Recognizes endogenous levels of Septin 5 protein
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human Septin 5. The exact sequence is proprietary.
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 39; Observed: 43 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	PNUTL1; Septin-5; Cell division control-related protein 1; CDCrel-1; Peanut-like protein 1
Gene Symbol	SEPT5
Entrez Gene	5413(Human); 18951(Mouse); 116728(Rat)
SwissProt	Q99719(Human); Q9Z2Q6(Mouse); Q9JJM9(Rat)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.


*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET


Septin 5 Rabbit Polyclonal Antibody

CAT. NO. APA14985


DATA



Western blot analysis of Septin 5 expression in Jurkat (A), NIH3T3 (B), mouse heart (C), mouse kidney (D), rat brain (E) whole cell lysates. (Predicted band size: 39; 42 kD; Observed band size: 43 kD)



Immunohistochemical analysis of Septin 5 staining in rat brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Septin 5 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AREX® Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.