

DATASHEET

MCEE Rabbit Polyclonal Antibody

CAT. NO. APA16114

KEY FEATURES

Target	MCEE	Source / Host	Rabbit
Reactivity	Human, Mouse	Clonality	Polyclonal
Applications	WB, IHC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

BACKGROUND

Methylmalonyl-CoA epimerase involved in propionyl-CoA metabolism.

APPLICATION

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:2000
IHC	1:50 - 1:200

*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

PRODUCT OVERVIEW


Description	Rabbit polyclonal antibody to MCEE
Specificity	Recognizes endogenous levels of MCEE protein
Antibody Type	Primary antibody
Immunogen	Recombinant fusion protein of human MCEE. The exact sequence is proprietary.
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 18 kD; Observed: 19 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	Methylmalonyl-CoA epimerase mitochondrial; DL-methylmalonyl-CoA racemase
Gene Symbol	MCEE
Entrez Gene	84693(Human); 73724(Mouse)
SwissProt	Q96PE7(Human); Q9D1I5(Mouse)

*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact info@arexbio.com or your local distributor.

*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

DATASHEET**MCEE Rabbit Polyclonal Antibody**

CAT. NO. APA16114

DATA

Western blot analysis of MCEE expression in HeLa (A), mouse brain (B) whole cell lysates. (Predicted band size: 18 kD; Observed band size: 19 kD)

STORAGE

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

NOTE

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.