

**DATASHEET**

**PARP16 Rabbit Polyclonal Antibody**

CAT. NO. APA18892

**KEY FEATURES**

Target	PARP16	Source / Host	Rabbit
Reactivity	Human	Clonality	Polyclonal
Applications	WB, IHC, FC	Conjugation	Unconjugated
Form / Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		Storage at -20°C

**BACKGROUND**

Intracellular mono-ADP-ribosyltransferase that plays a role in different processes, such as protein translation and unfolded protein response (UPR), through the mono-ADP-ribosylation of proteins involved in those processes, through the mono-ADP-ribosylation of proteins involved in those processes. Acts as an inhibitor of protein translation by catalyzing mono-ADP-ribosylation of ribosomal subunits, such as RPL14 and RPS6, thereby inhibiting polysome assembly and mRNA loading. Mono-ADP-ribosylation of ribosomal subunits is promoted by NMNAT2. Involved in the unfolded protein response (UPR) by ADP-ribosylating and activating EIF2AK3 and ERN1, two important UPR effectors. May also mediate mono-ADP-ribosylation of karyopherin KPNB1 a nuclear import factor.

**APPLICATION**

To ensure optimal assay performance, AREX recommends conducting reagent titration tailored to each testing system for optimal detection results.

WB	1:500 - 1:1000
IHC	1:50 - 1:200
FC	1:10 - 1:50

\*Results are sample-specific. Please refer to your local assay conditions and test parameters for reference.

**PRODUCT OVERVIEW**

Description	Rabbit polyclonal antibody to PARP16
Specificity	Recognizes endogenous levels of PARP16 protein.
Antibody Type	Primary antibody
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-terminal region of human PARP16. The exact sequence is proprietary.
Purification	The antibody was purified by immunogen affinity chromatography.
Molecular Weight	Predicted: 36 kD; Observed: 38 kD
Form/Buffer	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Alternative Names	C15orf30; Mono [ADP-ribose] polymerase PARP16; ADP-ribosyltransferase diphtheria toxin-like 15; ARTD15; Poly [ADP-ribose] polymerase 16; PARP-16
Gene Symbol	PARP16
Entrez Gene	54956(Human)
SwissProt	Q8N5Y8(Human)

\*AREX continuously optimizes our products. Webpage content may not reflect the latest updates. For inquiries, please contact [info@arexbio.com](mailto:info@arexbio.com) or your local distributor.


\*Clone Number, Reactivity, Source/Host and Clonality can be found in the product name and Key Features section above.

**DATASHEET**


**PARP16 Rabbit Polyclonal Antibody**

CAT. NO. APA18892


**DATA**



Western blot analysis of PARP16 expression in HepG2 (A), MDAMB435 (B) whole cell lysates. (Predicted band size: 36 kD; Observed band size: 38 kD)



Immunohistochemical analysis of PARP16 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Flow cytometric analysis of HepG2 cells using Anti-PARP16 Antibody. The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody at 37 °C for 60 min. The secondary antibody Goat Anti-Rabbit IgG (H&L) - AREX® Fluor 488 was incubated at 37 °C for 40 min. Isotype control antibody (blue line) was used under the same condition.

**STORAGE**

Store at 4°C short term. For long term storage, store at -20°C, avoiding freeze/thaw cycles.

**NOTE**

For Research Use Only. Not for diagnostic, therapeutics, prophylactic or in vivo use.